

Phenotypic and phylogenetic characterization of dominant culturable methanogens isolated from ricefield soils

C. Joulian^a, B. Ollivier^a, B.K.C. Patel^b, P.A. Roger^{a,*}

^a Laboratoire ORSTOM de Microbiologie des Anaérobies, Université de Provence, CESB/ESIL, Case 925, 163 Avenue de Luminy, 13288 Marseille cedex 9, France

^b School of Biomolecular and Biomedical Sciences, Faculty of Science and Technology, Griffith University, Brisbane, Qld. 4111, Australia

Received 26 June 1997; revised 16 October 1997; accepted 21 October 1997

Abstract

Populations of the four major trophic groups of methanogens were enumerated by most probable numbers (MPN) on selective media in a sample of 13 soils representative of major types of rice soils. Dominant strains were isolated and their phenotypic and phylogenetic characteristics were studied. MPN counts ranged from 10^2 to 10^6 g⁻¹ d.w. on H₂, from < 10 to 10^4 g⁻¹ d.w. on acetate, from < 10 to 10^5 on methanol, and from 50 to 10^6 on formate. In most soils, counts of hydrogenotrophs were higher than counts of acetotrophs, partly because acetotrophs were aggregated sarcinae difficult to separate into individual cells. Methylotrophs other than acetotrophic sarcinae were not recorded. In most soils, rods enumerated on formate were 5–400 times less abundant than those enumerated on H₂, indicating that hydrogenotrophic-non-formatotrophic rods are probably dominant in ricefields. Dominant strains isolated comprised: 15 hydrogenotrophic-non-formatotrophic rods affiliated to *Methanobacterium bryantii*; three hydrogenotrophic-formatotrophic rods affiliated to *Mb. formicum*; one hydrogenotrophic-formatotrophic rod not affiliated to a sequenced species; two sarcinae affiliated to *Methanosarcina barkeri* and *Methanosarcina mazei*; and one irregular coccus affiliated to *Methanoculleus marisnigri* – a species so far isolated from marine sediments only. Results from classical counts of methanogens and strains isolated from ricefields suggest the dominance of *Methanobacterium* spp. (mostly responsible for CH₄ production from H₂/CO₂) and *Methanosarcina* spp. (mostly responsible for CH₄ production from acetate) among culturable organisms. Both genera are probably ubiquitous. In particular, *Mb. bryantii* was isolated from 12 of the 13 soils. © 1998 Federation of European Microbiological Societies. Published by Elsevier Science B.V.

Keywords: Methane; Methanogen; MPN count; Ricefield; Soil; Phylogeny; *Methanobacterium*

1. Introduction

Methane (CH₄) has a high potential for absorbing infrared radiations and is therefore one of the major

gases involved in the greenhouse effect [1,2]. Up to 70–80% of atmospheric CH₄ is biogenic [3]. Waterlogged ricefields, because of anoxic conditions developing after flooding, are one of the major anthropogenic sources of CH₄ [4], however, the estimation of their contribution to the global CH₄ budget remains relatively imprecise [5].

Methane-producing bacteria are strict anaerobes

* Corresponding author. Tel.: +33 4 9182 8571; Fax: +33 4 9182 8570; E-mail: rogerpa@orstom.esil.univ-mrs.fr

Table 1
Primers used for amplification and sequencing of 16S rRNA gene of methanogens

| Primer | <i>E. coli</i> position | Sequence 5'-3' |
|-------------------------|-------------------------|--------------------------------|
| Forward | | |
| MS1 ^a | 406 | CCA CTC TTA ACG GGG TG |
| meth-ArFd3 ^b | 7 | G (A) TC CGT TTG ATC CTG GCG G |
| FARCH-9 ^b | 9 | CTG GTT GAT CCT GCC AG |
| F2 | 785 | CAG GAT TAG ATA CCC TGG TAG |
| F3 | 907 | AAA CTC AAA GGA ATT GAC GG |
| F4 | 1391 | TGT ACA CAC CGC CCG T |
| Reverse | | |
| MS2 ^a | 1342 | ACA CGC GAT TAC TAC GC |
| meth-ArchR1 | 337 | GCT GCG CCC CGT AGG GCC |
| R2 | 519 | GTA TTA CCG CGG CTG CTG |
| R4 | 907 | CCG TCA ATT CAT TTG AGT TT |
| R6 | 1494 | TAC GGT TAC CTT GTT ACG AC |
| Rd1 ^b | 1542 | AAG GAG GTG ATC CAG CC |

^aPrimers used for partial 16S rRNA gene amplification (about 940 bp).

^bPrimers used for complete 16S rRNA gene amplification and sequencing.

belonging to the *Archaea* domain. They are commonly isolated from natural anoxic environments, including freshwater and marine sediments, wet and waterlogged soils, the rumen, and the gut of insects [6–8]. They play an important role in these environments by performing the last step of anaerobic decomposition of organic matter, which is mineralized into CH₄ and CO₂. Radiotracer experiments showed that H₂ and acetate are the main energy sources used by methanogens in ricefields [9–11]. Those substrates result from fermentative metabolism or syntrophic associations degrading reduced compounds such as butyrate and propionate [12–14].

Despite numerous studies providing indirect evidence of the occurrence of methanogens in ricefields, few reports are available on their density [10,15–17], and species present in ricefields [11,17–20]. The only fully characterized species isolated from a ricefield are *Methanobrevibacter arboriphilicus* strain SA [21] and *Methanosarcina mazei* strain TMA [22]. Recently, a phylogenetic study has reported the presence of *Methanosarcina*, *Methanogenium*, *Methanosaeta*, and *Methanobacterium* in Japanese ricefield soils [23].

This paper reports on the study of the four major trophic groups of methanogens in ricefield soils. Populations were enumerated in a sample of 13 soils representative of major rice soils. Dominant strains were isolated and their phenotypic and phylogenetic characteristics were studied.

2. Materials and methods

2.1. Soils

Soils were collected from a range of ricefields in three countries (France, the Philippines, and USA), providing a representative sample with a broad range of physico-chemical properties: pH ranged from 4.5 to 7.8; organic C content ranged from 0.88 to 3.72%; organic N content ranged from 0.10 to 0.38%; available P ranged from 2 to 22 ppm. Soils were collected at the end of the crop cycle as composite samples. They were air-dried at ambient temperature as large clods and stored at room temperature. Before use, they were crushed and passed through a 5 mm sieve.

2.2. Reference bacterial strains

Methanobacterium bryantii (DSM 863), *Methanobacterium formicicum* (DSM 1535), *Methanobacterium wolfei* (DSM 2970), and *Methanobacterium thermoautotrophicum* (DSM 1053) were obtained from the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ), Braunschweig, Germany.

2.3. Cultivation and media

Hungate anaerobic techniques [24,25] were used

throughout this work. Basal medium contained (per liter): 1 g of NH_4Cl ; 0.3 g of KH_2PO_4 ; 0.3 g of K_2HPO_4 ; 0.2 g of $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$; 0.1 g of $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$; 0.1 g of KCl ; 0.5 g of CH_3COONa anhydrous; 0.6 g of NaCl ; 0.5 g of cysteine-HCl; 10 ml of trace element solution [26]; and 0.001 g of rezasurin. The pH of the medium was adjusted to 7 using a 10 M KOH solution. The medium was then boiled under a stream of O_2 -free N_2 and cooled at room temperature. It was distributed anaerobically into 60-ml serum bottles (20 ml/bottle) and into 20-ml Hungate tubes (5 ml/tube). Hungate tubes and serum bottles were flushed with N_2/CO_2 (80/20 v/v) and sterilized for 45 min at 110°C. After sterilization, 0.01 ml of 2% $\text{Na}_2\text{S}_2\text{O}_3$ and 0.05 ml of 10% NaHCO_3 (sterile, anaerobic solutions) were injected per ml of basal medium into the culture vessels.

Four selective media were prepared by adding to the sterilized basal medium, one of the four following substrates: formate (40 mM), methanol (40 mM), acetate (20 mM), which were added from sterile and anaerobic stock solution, or H_2/CO_2 (80/20 v/v, 2 bar), which was injected in the gas phase. For enumerations, these media were supplemented with 1 g l⁻¹ of yeast extract (Difco Laboratories, Detroit, MI) and 1 g l⁻¹ of bio-Trypticase (BioMérieux, Craponne, France). For preparing solid media, 1.6%

agar (Difco) was used. For growth requirement studies, vitamin (1% of Pfennig-Widdel [27] solution), various concentrations of yeast extract, and bio-Trypticase were added.

2.4. Counts of methanogens

The population density of the major trophic groups of methanogens were estimated by the most probable number (MPN) method (3 tubes per dilution). Successive 10-fold serial soil suspension dilutions were inoculated in the four selective media described above. Trophic groups of methanogens are detected, at least, on one of these media. Counts were duplicated by using two composite soil samples for each soil type. Methanogen growth was assayed by measuring CH_4 produced after 60 days of incubation at 37°C. Inoculated tubes containing medium supplemented with 1 g l⁻¹ of yeast extract and 1 g l⁻¹ of bio-Trypticase, where no substrate was added, served as control. A tube was considered positive when CH_4 produced was at least 5% higher than in the control. Populations were expressed as MPN per g dry soil. Because MPN counts have a low accuracy, counts (average of two replicates) were considered different if their ratio was higher than four [28].

Table 2
Counts of major trophic groups of methanogens

| Soil | Origin of soil | MPN of methanogens (g ⁻¹ soil d.w.) on | | | |
|--------------------------|-----------------|---|-------------------|-------------------|-------------------|
| | | H_2/CO_2 | Formate | Methanol | Acetate |
| Camargue | France | 9.6×10^3 | 1.8×10^3 | 2.5×10^3 | 5.0×10^3 |
| California 3 | USA | 1.0×10^2 | 1.3×10^2 | 2.8×10^2 | 1.8×10^2 |
| Amurao | The Philippines | 5.7×10^2 | 1.1×10^3 | 2.6×10^2 | 2.3×10^2 |
| Binalonan | The Philippines | 4.8×10^3 | 2.0×10^2 | 1.2×10^2 | 2.4×10^2 |
| Bugallon | The Philippines | 3.8×10^2 | 6.7×10^1 | 2.0×10^1 | 2.7×10^1 |
| Lal-lo | The Philippines | 2.1×10^2 | 5.3×10^1 | < 10 ¹ | 4.7×10^1 |
| Luisiana | The Philippines | 1.9×10^3 | 9.0×10^3 | 5.0×10^2 | 1.2×10^2 |
| Maahas | The Philippines | 7.0×10^4 | 1.0×10^3 | 2.9×10^2 | 1.3×10^2 |
| Maahas+salt ^a | The Philippines | 1.3×10^5 | 7.0×10^3 | 1.8×10^2 | 5.0×10^2 |
| Maahas+GM ^b | The Philippines | 5.0×10^4 | 1.5×10^3 | 1.8×10^2 | 5.0×10^2 |
| Maligaya | The Philippines | 1.8×10^4 | 4.3×10^2 | 2.2×10^2 | 1.8×10^2 |
| Pila | The Philippines | 3.6×10^4 | 9.0×10^1 | 4.0×10^3 | 2.9×10^3 |
| San Dionisio | The Philippines | 1.7×10^2 | 6.0×10^1 | 2.0×10^2 | < 10 ¹ |
| Tiaong | The Philippines | 2.3×10^6 | 1.6×10^6 | 1.9×10^5 | 2.2×10^4 |
| Urdaneta | The Philippines | 7.3×10^2 | 6.0×10^1 | 6.5×10^1 | 1.3×10^2 |

^a0.66 kg NaCl m⁻² applied before transplanting.

^bGreen manure (20 t ha⁻¹ fresh weight) incorporated before transplanting.

Table 3
Origin, morphology, and energy sources of isolated strains

| Soil | Strain | Form | Energy sources used for | | | | | |
|--------------------------|--------|---------|---------------------------------|---------------------------------|---------|----------|---------|----------|
| | | | Isolation | Growth | | | | |
| | | | | H ₂ /CO ₂ | Formate | Methanol | Acetate | Alcohols |
| Camargue | RiH2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| | FCam | Rod | Formate | + | + | – | – | 2-P |
| | CoCam | Coccoid | Formate | + | + | – | – | – |
| | Sar | Sarcina | Methanol | + | – | + | + | – |
| California 3 | H2Cal3 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| Amurao | TAH2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| Binalonan | PBH2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| Bugallon | PDH2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| Lal-lo | CBH2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| Luisiana | LuiH2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| Maahas | MaH2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| | H2Sol2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| | FSol1 | Rod | Formate | + | + | – | – | – |
| Maahas+salt ^a | SelH2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| Maahas+GM ^b | GMH2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| Maligaya | MaliH2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| Pila | PiH2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| | FPI | Rod | Formate | + | + | – | – | – |
| | SarPi | Sarcina | Methanol | + | – | + | + | B |
| San Dionisio | IAH2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |
| Tiaong | TiaH2 | Rod | H ₂ /CO ₂ | + | + | – | – | – |
| Urdaneta | PAH2 | Rod | H ₂ /CO ₂ | + | – | – | – | 2-P |

^{a,b}As in Table 2.

2-P: 2-propanol; B: butanol.

2.5. Enrichment and isolation procedure

Pure cultures of methanogens were obtained from the last positive tubes, by three repeated applications of the agar shake dilution method [24]. For final purification, single colonies were picked up, using an anaerobic glove box with a N₂/H₂ atmosphere (95/5 v/v). Colonies were diluted in liquid media and the last positive dilution was checked for purity by microscopic examination and by verifying the absence of growth in a medium containing 1 g l⁻¹ bio-Trypticase and 1 g l⁻¹ yeast extract, to which was added either (1) 20 mM glucose (to detect fermentative bacteria) or (2) 20 mM sulfate and 20 mM lactate (to detect sulfate-reducing bacteria).

2.6. Morphology and growth characteristics

Growth characteristics of the isolates were determined in duplicate Hungate tubes. Media were enriched with 1 g l⁻¹ of yeast extract and, when neces-

sary, 1 g l⁻¹ of bio-Trypticase. The temperature range for growth was tested from 15°C to 50°C in water baths. Optimum NaCl concentration was tested at 37°C using concentrations ranging from 0 to 60 g l⁻¹. The pH range for growth was determined by adding various concentration of 10% NaHCO₃, 8% Na₂CO₃, or 0.5 M HCl solutions to the culture medium, to obtain pHs ranging from 4 to 9. The following substrates were tested for growth: 2 bar H₂/CO₂, 40 mM formate, 40 mM methanol, 20 mM acetate, 10 mM 1-propanol, 10 mM 2-propanol, 10 mM 1-butanol, and 10 mM 2-butanol.

2.7. Analytical techniques

Bacterial cultures were observed with a Nikon Optiphot microscope equipped with a Nikon FX35 camera and a Zeiss, standard 20, microscope equipped with epifluorescence. Bacterial growth was quantified (1) with a Shimadzu UV 160 A spectrophotometer by measuring turbidity increase at




Fig. 1. Restriction profiles of the isolated rods. *Mb. bryantii* group included strains RiH2, H2Ca3, TAH2, PBH2, PDH2, CBH2, LuiH2, MaH2, H2So2, SelH2, GMH2, MaliH2, PAH2, IAH2 and PiH2; *Mb. formicicum* group included strains FCam, FSol1 and TiaH2. L, ladder; a, *Bam*HI; b, *Cfo*I; c, *Sau*3A; d, *Taq*I.

580 nm directly in Hungate tubes, or (2) through CH_4 production for aggregated forms such as members of the genus *Methanosarcina* or methanogenic cultures with low density. Methane was measured with a Girdel series 30 flame ionization detector gas chromatograph (column SP 1000 with chromosorb WAW and 1% H_3PO_4) operated at 150°C with N_2 as carrier gas.

2.8. Phylogenetic analysis

2.8.1. Restriction profile

Part of the 16S rRNA gene (about 940 bp) of the isolated *Methanobacterium* strains was amplified. The sequences of forward (MS1), and reverse (MS2) primers used were determined by comparing the 16S rRNA sequences of *Methanobacterium* species available in the GenBank database. Primer sequences are described in Table 1. Primers were synthesized by Bioprobe® systems (Montreuil, France). PCR was performed directly from 1 μl of culture. The PCR product was digested with four restriction enzymes, *Bam*HI, *Cfo*I, *Sau*3A and *Taq*I (Boehringer Mannheim, Germany).

2.8.2. 16S rRNA gene sequencing

Semi-purified DNA was extracted for amplification of the 16S rRNA gene using the following protocol. Pure cultures (50 ml) were centrifuged for 20 min at $10\,000\times g$ and pellets were resuspended in 50 μl of lysis buffer (50 mM Tris-HCl, pH 7.2; 50 mM EDTA; 3% sodium dodecyl sulfate). Suspensions were microwaved, with lids open, at 600 W (15 s

on, 5 s off; repeated four times), and were then placed at 80°C for 2 h. This treatment was repeated three times. Then, 350 μl of lysis buffer was added and the suspensions were placed at 80°C. After overnight incubation, suspensions were mixed vigorously with 400 μl of phenol:chloroform:isoamyl alcohol (25:24:1) and centrifuged for 15 min at $10\,000\times g$. The top aqueous phase was removed carefully to avoid collecting cell material from the interface. Then, 10 μl of cold isopropanol and 5 μl of 5 M sodium acetate (pH 5.8) were added to the aqueous phase. The suspensions were centrifuged, the nucleic acid pellets were washed with 500 μl of 70% cold ethanol and dried in a desiccator. Pellets were resuspended in 50 μl of TE buffer (10 mM Tris-HCl; 1 mM EDTA) and stored at -20°C until use.

The sequences of primers used for amplification and sequencing are described in Table 1. The 16S rRNA gene was amplified as described by Love et al. [29] and Redburn and Patel [30] using Farch9 and Rd1 as PCR primers. In addition, a new forward primer, meth-ArFd3, specific to *Methanobacterium* strains, was used for amplification. The purified PCR product was sequenced directly with an ABI automated DNA sequencer, using a Prism dideoxy terminator cycle sequencing kit, as recommended by the manufacturer (Applied Biosystems, Ltd., Foster City, CA). Primers used were described by Redburn and Patel [30]. In addition, a new reverse primer specific to *Methanobacterium* strains, meth-ArchR1, was used to sequence the beginning of the gene. New primers were synthesized by CMCB (University of the Southern Cross, Lismore, Australia).

The 16S rRNA sequences of the isolated strains and 16S rRNA sequences of various members of the *Archaea* domain, obtained from the RNA Database Project and from the GenBank database, were manually aligned using the sequence editor ae2 [31]. Positions at which the sequences and/or alignment were uncertain were omitted from the analysis, and pairwise evolutionary distances for unambiguous nucleotides were computed by the method of Jukes and Cantor [32]. Dendrograms were constructed from these distances by the neighbor-joining method. All programs are available as part of the PHYLIP package [33].

3. Results

3.1. Estimation of methanogenic populations

Population of methanogens were quantified in the 13 ricefield soils using four substrates (Table 2). MPN counts ranged from 10^2 to 2×10^6 g⁻¹ d.w. for hydrogenotrophs, from 50 to 2×10^6 g⁻¹ d.w. for formatotrophs, from <10 to 2×10^5 g⁻¹ d.w. for methylotrophs, and from <10 to 2×10^4 g⁻¹ d.w. for acetotrophs.

Hydrogenotrophs were dominant in eight of the 13 soils, being 5–400 times more abundant than formatotrophs, 9–650 times more abundant than methylotrophs, and 5–538 times more abundant than acetotrophs. In Tiaong soil, both hydrogenotrophs and formatotrophs were dominant; their populations were similar and exhibited the highest density among the soils tested. In California 3 and Amurao soils, the four methanogenic trophic groups exhibited approximately the same densities within each soil.

When methanol or acetate was used as energy source, *Methanosarcina*-like forms were dominant and population densities were of the same order on both substrates in most of the soils. When H₂ or formate was used as energy source, rods were dominant. They showed the typical fluorescence of methanogens under UV light and phenotypic characteristics similar to members of the *Methanobacterium* genus. Population densities were usually higher when enumerated on H₂/CO₂.




Fig. 2. Dendrogram showing the position of the methanogenic strains isolated from ricefields among representative methanogens of various origins.

3.2. Isolation

The dominant morphotypes isolated on H₂ or formate were rods, which were observed as single cells or chains, depending on the growth phase. In Camargue and Pila soils, irregular coccoid cells were also present in tubes containing formate as energy source, but rods still dominated. Enrichments on acetate or methanol led to the development of sarcina-like morphotypes. Their isolation was further performed on methanol. In addition, in some enrichments we observed the presence of spirilla showing a fluorescence typical of methanogens under UV microscopic examination. Attempts to isolate these spirilla were unsuccessful.

Twenty-two methanogenic strains were isolated from the 13 soils studied (Table 3). We found 19 rods, phenotypically related to the genus *Methanobacterium*, among which 15 were hydrogenotrophs unable to utilize formate, and four used both H₂ and formate as energy sources. *Methanosarcina*-like strains were present in all soils; two strains were isolated. An irregular methanogenic coccus was observed in tubes containing formate in Camargue and Pila soils; one strain was isolated.

3.3. Phylogenetic and phenotypic characterization

As isolated *Methanobacterium* strains were too numerous to be all studied for their phenotypic characteristics and sequenced, we first characterized them from their RFLP profiles using four restriction enzymes (*Bam*HI, *Cfo*I, *Sau*3A, and *Taq*I) and about 940 bp of the 16S rRNA gene. We compared their profiles with those of the four *Methanobacterium* species with known 16S rRNA gene sequences (*Mb. bryantii*, *Mb. formicicum*, *Mb. wolfei*, and *Mb. thermoautotrophicum*). These four species exhibited four different profiles. The 19 isolated rods exhibited three profiles (Fig. 1), suggesting that three species of *Methanobacterium* were present. The 15 hydrogenotrophic-non-formatotrophic rods had a profile similar to that of *Mb. bryantii*, and three of the four formatotrophic-hydrogenotrophic rods had a profile similar to that of *Mb. formicicum*. A profile different from the four reference profiles was obtained for one formatotrophic-hydrogenotrophic rod (strain FPi). Then, we sequenced the complete

16S rRNA gene of one strain of each group (strain RiH2 for *Mb. bryantii* group, strain FCam for *Mb. formicicum* group, and strain FPi for the unknown *Methanobacterium* sp. group), using 10 primers (Table 1). The analysis of the sequences confirmed the determination of the three species of *Methanobacterium* isolated. Finally, we used the other isolated strains of *Methanobacterium* to sequence the most variable region of their 16S rRNA gene, located at the beginning of the gene, using primers meth-ArFd3 and meth-ArchR1 (Table 1). Results confirmed the groups established with the restriction profiles.

These studies showed that the 15 hydrogenotrophic-non-formatotrophic rods were related to the species *Mb. bryantii*. Three of the four formatotrophic rods were related to *Mb. formicicum*. The formatotrophic rod, strain FPi, was significantly distant from currently sequenced *Methanobacterium* species.

To determine the phylogeny of all isolated methanogens, we sequenced the complete or almost complete 16S rRNA gene, using 10 primers (Table 1), of the six different species isolated: the three rods, as

Table 5
Major characteristics of the isolated strains and related species^a

| Microorganism | Culture conditions | | | | | | | | ARNr 16S phylogeny | |
|--------------------------------|---------------------|---------------------|-----------------|---------------------------------|---------|----------|---------|----------|------------------------------|----------------|
| | Optimum (and range) | | | Substrate for growth | | | | | Most closely related species | Similarity (%) |
| | Temperature (°C) | Salinity (g/l) | pH | H ₂ /CO ₂ | Formate | Methanol | Acetate | Alcohols | | |
| <i>Methanobacterium</i> | | | | | | | | | | |
| <i>bryantii</i> ^b | 37–39 | n.d. | 6.9–7.2 | + | – | – | – | + | | |
| <i>formicicum</i> ^b | 37–45 | n.d. | 6.6–7.8 | + | + | – | – | ± | | |
| RiH2 (Camargue) ^c | 35 (25–42) | 0–25 | 6.8 (6.0–8.5) | – | + | – | – | 2-P | <i>Mb. bryantii</i> | 99.2 |
| FCam (Camargue) ^c | 40 (25–42) | 5 (0–30) | 7–7.2 (6.0–8.5) | + | + | – | – | 2-P | <i>Mb. formicicum</i> | 97.9 |
| FPi (Pila) ^c | 40 (20–42) | 5 (0–30) | 7 (6.0–8.0) | + | + | – | – | – | <i>Mb. bryantii</i> | 96.5 |
| <i>Methanosarcina</i> | | | | | | | | | | |
| <i>barkeri</i> ^b | 30–40 | n.d. | neutral | + | – | + | + | – | | |
| <i>mazei</i> ^b | 30–40 | n.d. | 7–7.2 | + | – | + | + | n.d. | | |
| Sar (Camargue) ^c | 35–37 (25–42) | 0 (0–10) | (5.5–7.5) | + | – | + | + | – | <i>Ms. barkeri</i> | 99.0 |
| SarPi (Pila) ^c | 40 (20–42) | (0–60) ^d | 7.5 (6.0–8.5) | + | – | + | + | B | <i>Ms. mazei</i> | 99.8 |
| <i>Methanoculleus</i> | | | | | | | | | | |
| <i>marisnigri</i> ^b | 20–25 | 60 | 6.8–7.3 | + | + | – | – | + | | |
| CoCam (Camargue) ^c | 37 (20–42) | (0–60) ^d | 7 (6.8–8.2) | + | + | – | – | – | <i>Mc. marisnigri</i> | 98.4 |

^aCharacteristics of related species according to Garcia [8].

^bReference strains.

^cSoil of origin.

^dConcentrations > 60 g l⁻¹ of NaCl not tested.

2-P: 2-propanol; B: 1-butanol.

n.d.: not determined.

described above (strains RiH2, FCam and FPi), the two sarcinae, and the irregular coccus. Sequences were aligned with the sequences of representative methanogens, and phylogenetic analysis was performed from the Jukes-Cantor similarity matrix [32] (Table 4). A dendrogram was obtained from this matrix by the FITCH method [33] (Fig. 2). Results indicated that five of the six isolated species were related phylogenetically but also phenotypically to known species (Table 5). RiH2 (sequence accession number AF028688) belongs to *Methanobacterium bryantii* (M59124) (average similarity of 99.2%), FCam (AF028689) to *Methanobacterium formicicum* (M36508) (average similarity of 97.9%). SarPi (AF028691) belongs to *Methanosarcina mazei* (U20151) (average similarity of 99.8%), and Sar (AF028692) to *Methanosarcina barkeri* (M59144) (average similarity of 99%). The irregular coccus, strain CoCam (AF028693), which uses H₂ and formate, belongs to *Methanoculleus marisnigri* (M59134) (average similarity of 98.4%). Strain FPi (AF028690) showed marked phylogenetic differences from its closest relatives, *Methanobacterium bryantii* (average similarity of 96.5%) and *Methanobacterium formicicum* (average similarity of 94.7%). In contrast to its phylogenetically closest relative, *Mb. bryantii*, it used formate as energy source (Table 5).

4. Discussion

We enumerated methanogens in dry soil samples originating from 13 ricefields representing a broad range of physico-chemical properties and geographical origins. Since acetate and H₂/CO₂ were demonstrated to be the major methanogenic substrates in wetland ricefields [9–11], enumerations were performed on selective media containing each of these energy sources. In addition, we also used two additional selective media containing (1) methanol, to record strict (non-acetoclastic) methylotrophs and (2) formate, since hydrogenotrophic methanogens do not necessarily use formate [8]. Formate might be important in the anaerobic degradation of organic matter, because, like H₂, it is involved in the interspecies transfer during oxidation of reduced compounds [14,34].

MPN counts of methanogens on H₂ ranged from

10² to 10⁶ g⁻¹ d.w.; counts on acetate ranged from < 10 to 10⁴ g⁻¹ d.w. Populations were in a range similar to those reported in 29 Senegalese ricefields by Garcia et al. [15] (10²–10⁷ g⁻¹ d.w.). In an Italian ricefield, Schütz et al. [10] and Mayer and Conrad [16] counted 10⁴–10⁵ acetotrophs g⁻¹ d.w. and 10⁵–10⁶ hydrogenotrophs g⁻¹ d.w.

In most soils, counts of hydrogenotrophs were higher than counts of acetotrophs. However, the lower counts of acetotrophs mostly resulted from a methodological bias, hydrogenotrophs being mostly rods, whereas acetotrophs were mostly sarcinae. Sarcinae are known to develop as dense aggregates, difficult to separate into individual cells, thus their populations are underestimated by MPN counts, which mostly record the number of aggregates [20].

In most soils, counts on methanol did not show significant differences with counts on acetate. Sarcinae dominated on both substrates; non-acetoclastic-methylotrophic methanogens, such as members of the genera *Methanococcoides* or *Methanobolus*, were not evidenced. These results suggest that methanogenesis from methanol released from the fermentation of pectin [35] originating from the photosynthetic aquatic biomass (aquatic macrophytes, micro- and macro-algae, and cyanobacteria) is probably mainly due to the activity of *Methanosarcina*.

In most soils, rods enumerated on formate were 5–400 times less abundant than those enumerated on H₂, indicating that hydrogenotrophic-non-formatotrophic rods were dominant in the ricefield soils tested, since all formatotrophs also use H₂/CO₂. In all soils, dominant hydrogenotrophic methanogens belonged to the genus *Methanobacterium*.

Twenty-two strains were isolated from the highest dilution tubes. They included: two sarcinae isolated on methanol, one irregular coccus isolated on formate, 16 hydrogenotrophic rods isolated on H₂/CO₂, and three formatotrophic rods isolated on formate. Among the hydrogenotrophic rods, one was found to be also a formatotroph (Table 3).

On the basis of phenotypic studies, all the isolated rods were affiliated to the genus *Methanobacterium*. Fifteen strains were related to *Mb. bryantii* and three to *Mb. formicicum*. One strain (FPi) presented sufficient phenotypic and phylogenetic differences from its closest relative, *Mb. bryantii* (average similarity

96.5%), to consider it a new species (work in progress).

Surprisingly, an irregular methanogenic coccus, phylogenetically related to *Methanoculleus marisnigri* (only isolated so far from marine sediments [36]) was isolated from the Camargue soil. This soil is of estuarine origin, but cannot be considered saline (exchangeable Na: 0.9 mEq/100 g). Members of the genus *Methanoculleus* are halotolerant microorganisms [37] and *M. marisnigri* in particular might be adapted to a wide range of NaCl concentrations. This probably explains its occurrence in marine but also in non-saline environments such as ricefields.

The isolated sarcinae were related to the species *Methanosarcina barkeri* and *Methanosarcina mazei*, which have already been isolated from a broad range of environments, including sediments and digesters. The latter species was also recently isolated from a Japanese ricefield [22].

Our results seem to indicate that, in ricefields, *Methanobacterium* spp. are mostly responsible for CH₄ production from H₂/CO₂ and *Methanosarcina* spp. for CH₄ production from acetate. Both genera are probably ubiquitous. In particular, *Mb. bryantii* was isolated from 12 of the 13 soils. However, beside members of these two genera, we have isolated a member of the genus *Methanoculleus*.

Although results from classical isolation of methanogens from ricefields suggest the ubiquity and dominance of *Methanobacterium* spp. among culturable organisms, Kudo et al. [23], using PCR amplification of archaeobacterial 16S ribosomal DNA from extracted soil DNA in nine Japanese ricefield soils, reported the presence of *Methanobacterium* in only one soil, where it was not dominant. On the other hand, similar to our results, this Japanese study also reported the presence of members of the genera *Methanosarcina* – including *Msr. mazei* and *Msr. barkeri* – and *Methanogenium* or *Methanoculleus*. From their results, Kudo et al. [23] concluded to the dominance of *Methanosarcina* in five soil samples, to that of *Methanogenium* in two soils and *Methanosaeta* in two soils.

The somewhat conflicting results regarding *Methanobacterium* occurrence may indicate that this genus is absent in most Japanese ricefield soils, whereas it is widely represented in soils of the Philippines, USA and France. However, these results might also be

due to bias inherent in both methods. On the one hand the DNA extraction and 16S rDNA clone analysis methods used by Kudo et al. [23] possibly do not fully reflect the diversity of the methanogenic flora. On the other hand the MPN method we used is possibly selective for *Methanobacterium* and might lead to the erroneous conclusion of its dominance when it is only present.

At this moment it is difficult to say which technique yield the most reliable results in terms of presence and relative abundance of genera of methanogens.

All together, these results indicate that methanogen biodiversity in ricefields is not yet elucidated. It clearly demonstrates the need for simultaneous classical microbiological studies and in situ phylogenetic analysis to assess microbial populations in soils.

Acknowledgments

The authors thank J.-L. Garcia for his helpful comments on the manuscript.

References

- [1] Wang, W.C., Yung, Y.L., Lacin, A.A., Mo, T. and Hansen, J.E. (1976) Greenhouse effects due to man-made perturbations of trace gases. *Science* 194, 685–690.
- [2] Lelieveld, J., Crutzen, P.J. and Brühl, C. (1993) Climate effect of atmospheric methane. *Chemosphere* 26, 739–768.
- [3] Wahlen, M., Tanaka, N., Henry, R., Deck, B., Zeglen, J., Vogel, J.S., Southon, J., Shemesh, A., Fairbanks, R. and Broecker, W. (1989) Carbon-14 in methane sources and in atmospheric methane: the contribution from fossil carbon. *Science* 245, 286–290.
- [4] Minami, K., Mosier, A. and Sass, R. (Eds.) (1994) CH₄ and NO₂ Global Emission and Controls from Ricefields and Other Agricultural and Industrial Sources. NIAES Series 2. Yokendo, Tokyo.
- [5] Sass, R.L. (1994) Short summary chapter for methane. In: CH₄ and NO₂ Global Emission and Controls from Ricefields and Other Agricultural and Industrial Sources (Minami, K., Mosier, A. and Sass, R., Eds.), pp. 1–7. NIAES Series 2. Yokendo, Tokyo.
- [6] Mah, R.A. and Smith, M.R. (1981) The methanogenic bacteria. In: *The Prokaryotes*, Vol. 1 (Starr, M.P., Stolp, H., Trüper, H.G., Balows, A. and Schlegel H.G., Eds.), pp. 948–977. Springer-Verlag, Berlin.
- [7] Bowmann, A.F. (Ed.) (1989) *Soils and the Greenhouse Effect*. Wiley, New York.

- [8] Garcia, J.-L. (1990) Taxonomy and ecology of methanogens. FEMS Microbiol. Rev. 87, 297–308.
- [9] Takai, Y. (1970) The mechanism of methane fermentation in flooded paddy soil. Soil Sci. Plant Nutr. 16, 238–244.
- [10] Schütz, H., Seiler, W. and Conrad, R. (1989) Processes involved in formation and emission of methane in rice paddies. Biogeochemistry 7, 33–53.
- [11] Conrad, R., Bak, F., Seitz, H.J., Thebrath, B., Mayer, H.P. and Schütz, H. (1989) Hydrogen turnover by psychrotrophic homoacetogenic and mesophilic methanogenic bacteria in anoxic paddy soil and lake sediment. FEMS Microbiol. Ecol. 62, 285–294.
- [12] Schink, B. (1992) Syntrophism among prokaryotes. In: The Prokaryotes, 2nd edn. (Balows, A., Trüper, H.G., Dworkin, M., Harder, W. and Schleifer, K.-H., Eds.), pp. 276–299. Springer Verlag, New York.
- [13] Stams, A.J.M. (1994) Metabolic interactions between anaerobic bacteria in methanogenic environments. Antonie van Leeuwenhoek 66, 271–294.
- [14] Dong, X. and Stams, A.J.M. (1995) Evidence for H₂ and formate formation during syntrophic butyrate and propionate degradation. Anaerobe 1, 35–39.
- [15] Garcia, J.-L., Raimbault, M., Jacq, V., Rinaudo, G. and Roger, P.A. (1974) Activités microbiennes dans les sols de rizières du Sénégal: relations avec les propriétés physico-chimiques et influence de la rhizosphère. Rev. Ecol. Biol. Sol 11, 169–185.
- [16] Mayer, H.P. and Conrad, R. (1990) Factors influencing the population of methanogenic bacteria and the initiation of methane production upon flooding of paddy soil. FEMS Microbiol. Ecol. 73, 103–112.
- [17] Joulain, C., Ollivier, B., Neue, H.-U. and Roger, P.A. (1996) Microbiological aspects of methane emission by a ricefield soil from the Camargue (France): 1. Methanogenesis and related microflora. Eur. J. Soil Biol. 32, 71–80.
- [18] Raimbault, M. (1981) Inhibition de la formation de méthane par l'acétylène chez *Methanosarcina barkeri*. Cah. ORSTOM, Sér. Biol. 43, 45–51.
- [19] Rajagopal, B.S., Belay, N. and Daniels, L. (1988) Isolation and characterization of methanogenic bacteria from rice paddies. FEMS Microbiol. Ecol. 53, 153–158.
- [20] Fetzer, S., Bak, F. and Conrad, R. (1993) Sensitivity of methanogenic bacteria from paddy soil to oxygen and desiccation. FEMS Microbiol. Ecol. 12, 107–115.
- [21] Asakawa, S., Morii, H., Akagawa-Matsushita, M., Koga, Y. and Hayano, K. (1993) Characterization of *Methanobrevibacter arboriphilicus* SA isolated from a paddy field soil and DNA-DNA hybridization among *M. arboriphilicus* strains. Int. J. Syst. Bacteriol. 43, 683–686.
- [22] Asakawa, S., Akagawa-Matsushita, M., Morii, H., Koga, Y. and Hayano, K. (1995) Characterization of *Methanosarcina mazei* TMA isolated from a paddy field soil. Curr. Microbiol. 31, 34–38.
- [23] Kudo, Y., Nakajima, T., Miyaki, T. and Oyaizu, H. (1997) Methanogen flora of paddy soils in Japan. FEMS Microbiol. Ecol. 22, 39–48.
- [24] Hungate, R.E. (1969) A roll tube method for cultivation of strict anaerobes. In: Methods in Microbiology, Vol. 3B (Norris, J.R. and Ribbons, D.W., Eds.), pp. 117–132. Academic Press, London.
- [25] Macy, J.M., Snellen, J.E. and Hungate, R.E. (1972) Use of syringe methods for anaerobiosis. Am. J. Clin. Nutr. 25, 1318–1323.
- [26] Balch, W.E., Fox, G.E., Magrum, L.J., Woese, C.R. and Wolfe, R.S. (1979) Methanogens, reevaluation of a unique biological group. Microbiol. Rev. 43, 260–296.
- [27] Pfennig, N., Widdel, F. and Trüper, H.G. (1981) Dissimilatory sulfate-reducing bacteria. In: The Prokaryotes, Vol. 1 (Starr, M.P., Stolp, H., Trüper, H.G., Balows, A. and Schlegel, H.G., Eds.), pp. 926–940. Springer-Verlag, Berlin.
- [28] Roger, P.A., Jimenez, R. and Santiago-Ardales, S. (1991) Methods for studying blue-green algae in ricefields: distributional ecology of the organisms, sampling strategies and estimation of abundance. Int. Rice Res. Paper Series No. 150. The International Rice Research Institute, P.O. Box 933, Manila.
- [29] Love, C.A., Patel, B.K.C., Nichols, P.D. and Stackebrandt, E. (1993) *Desulfotomaculum australicum* sp. nov., a thermophilic sulfate-reducing bacterium isolated from the Great Artesian Basin of Australia. Syst. Appl. Microbiol. 16, 244–251.
- [30] Redburn, A.C. and Patel, B.K.C. (1993) Phylogenetic analysis of *Desulfotomaculum thermobenzoicum* using polymerase chain reaction-amplified 16S rRNA-specific DNA. FEMS Microbiol. Lett. 113, 81–86.
- [31] Maidak, B.L., Olsen, G.J., Larsen, N., Overbeek, R., McCaughey, M.J. and Woese, C.R. (1996) The ribosomal database project (RDP). Nucleic Acids Res. 24, 82–85.
- [32] Jukes, T.H. and Cantor, C.R. (1969) Evolution of protein molecules. In: Mammalian Protein Metabolism (Munro, H.N., Ed.), pp. 21–32. Academic Press, New York.
- [33] Felsenstein, J. (1993) PHYLIP (Phylogenetic Inference Package) version 3.51c. Distributed by the author. Department of Genetics, University of Washington, Seattle, WA.
- [34] Thiele, J.H. and Zeikus, J.G. (1988) Control of interspecies electron flow during anaerobic digestion: significance of formate transfer versus hydrogen transfer during syntrophic methanogenesis in flocs. Appl. Environ. Microbiol. 54, 20–29.
- [35] Schink, B. and Zeikus, J.G. (1980) Microbial methanol formation: a major end product of pectin metabolism. Curr. Microbiol. 4, 387–389.
- [36] Romesser, J.A., Wolfe, R.S., Mayer, F., Spiess, E. and Walther-Mauruschat, A. (1979) *Methanogenium*, a new genus of marine methanogenic bacteria, and characterization of *Methanogenium cariaci* sp. nov. and *Methanogenium marisnigri* sp. nov. Arch. Microbiol. 121, 147–153.
- [37] Maestrojuan, G.M., Boone, D.R., Xun, L., Mah, R.A. and Zhang, L. (1990) Transfer of *Methanogenium bourgense*, *Methanogenium marisnigri*, *Methanogenium olentangyi*, and *Methanogenium thermophilicum* to the genus *Methanoculleus* gen. nov., emendation of *Methanoculleus marisnigri* and *Methanogenium*, and description of new strains of *Methanoculleus bourgense* and *Methanoculleus marisnigri*. Int. J. Syst. Bacteriol. 40, 117–122.